

Anti-Inflammatory

Highly Effective Topical Anti-Inflammatory and Pain Reliever without All the Systemic Effects

Medications that relieve inflammation and pain and relax muscles can cause undesirable side effects, including cardio-vascular complications, drowsiness, gastric irritation and unwanted systemic side effects. Topical administration (e.g., cream, gel and ointment) , on the other hand, delivers the drug directly to the disease site and produces high local drug concentration, while reducing unwanted side effects.



Ketoprofen (Orudis) and/or Piroxicam (Feldene) topical creams have anti-inflammatory and analgesic effects and are used in the local and symptomatic treatment of:

- Acute and chronic rheumatoid arthritis
- Tenosynovitis (e.g., Tendonitis)
- Acute and chronic osteo and peri arthritis
- Gonarthroses (inflammation of the knee joint)
- Bursitis (e.g., Tennis Elbow)
- Carpal Tunnel Syndrome
- Contusions
- Dislocations
- Edema
- Sprains
- Myalgias
- Myositis
- Ankylosing spondylitis (inflammation of the spine joints)
- Phlebitis
- Hematoma
- Superficial Thrombosis
- Acute musculoskeletal disorder
- Soft tissue injuries (e.g., ankle sprains, etc.)

These creams are often prescribed by Arthritis Specialists, Orthopedic Surgeons, Sports Medicine Specialists and Rehabilitation Therapists.

With a prescription from a physician, we at The Medicine Shoppe can prepare Ketoprofen with Piroxicam for additional anti-inflammatory effects and Ketoprofen with Lidocaine for anti-inflammatory and numbing effects. Ketoprofen and Piroxicam are both non-steroidal anti-inflammatory drugs (NSAIDs). Ketoprofen is structurally related to Fenoprofen, Ibuprofen and Naproxen. Piroxicam is structurally unrelated to other NSAIDs. Gabapentin (Neurontin), Ketamine (ketalar) , Misoprostol (Cytotec), Capsaicin, Amitriptyline (Elavil), Baclofen and many other active drugs could be added to the above for neurological (e.g., shingles and diabetic neuropathy) and other types of pain relief.

BASE USED

These creams are prepared in a transdermal pain base (PLO cream) vehicle for enhanced penetration.

DIRECTIONS

Apply the compound to the affected area, in a thin smooth film, 3 to 4 times daily (as needed). If wearing a brace over the cream, make sure to first air out the area for approximately 30 minutes. This cream is easily removed with water.

HOW TO ORDER

These compounds can be called into The Medicine Shoppe by a physician and, we can compound them for you. Prescription details could be as such:

- Ketoprofen 10% or 20%, in TD cream
- Ketoprofen 15%, Piroxicam 2%, in TD cream
- Ketoprofen 15%, Lidocaine 5%, in TD cream
- Ketoprofen 10%, Ibuprofen 10%, Piroxicam 2% in TD cream
- Diclofenac 2 - 5%, in TD cream
- Ibuprofen 20%, in TD cream
- Ibuprofen 20% / Cyclobenzaprine 1%, in TD cream
- Ibuprofen 20% / Cyclobenzaprine 3%, in TD cream

There are other variations that can be used and we would be glad to discuss them with your physician. Contact us today at **585-396-9970** to learn more.